Name……

Address……

……

……

CHI……

Affix patient label

IV Morphine Administration Proforma <10kg

Date: **Time:**

Indication for morphine administration:

Allergies?

Weight……..kg

Analgesia received:

Does the patient need further analgesia? Prescribe on ED card as required

Use Morphine 10mg/ml injection and Sodium Chloride 0.9%. Add 0.4ml (4mg) of Morphine 10mg/1ml to 19.6ml of 0.9% Sodium Chloride to make a total of 20ml. This gives a 200 microgram in 1ml solution. Exact dose (see below) should be drawn from syringe, labelled and taken to the bedside. Dose and flush should be given as a slow bolus over 5 minutes.

***If the patient weighs less than the half kg increment please round dose down e.g. if weight is 3.4 kg round down to dose for 3kg**

Weight in kg	Max dose in micrograms	Volume in mls of 200mcg/ml solution
2.5	125mcg (50mcg/kg)	0.62mls
3.0	150mcg (50mcg/kg)	0.75mls
3.5	175mcg (50mcg/kg)	0.87mls
4.0	200mcg (50mcg/kg)	1ml
4.5	225mcg (50mcg/kg)	1.10mls
5.0	500mcg (100mcg/kg)	2.50mls
5.5	550mcg (100mcg/kg)	2.75mls
6.0	600mcg (100mcg/kg)	3mls
6.5	650mcg (100mcg/kg)	3.25mls
7.0	700mcg (100mcg/kg)	3.50mls
7.5	750mcg (100mcg/kg)	3.75mls
8.0	800mcg (100mcg/kg)	4mls
8.5	850mcg (100mcg/kg)	4.25mls
9.0	900mcg (100mcg/kg)	4.50mls
9.5	950mcg (100mcg/kg)	4.75mls

All drugs should be prescribed on the ED card

Adjunctive Agents & Indications; Apnoea **NALOXONE** (100mcg/kg IV bolus) Repeat at intervals of 1 minute to a total max dose 2mg Calculated Dose:……. Max Dose:……

Pre- Administration checklist

Emergency equipment available: Airway trolley ☐ Suction ☐ Bag valve mask ☐ Non-rebreathe mask ☐	
Continuous SpO2 monitoring in situ	
Full set of observations recorded before administration	
Morphine prescribed on ED card and prepared	
Clinician prescribing /administering the medication has overseen the morphine preparation Name:	
Naloxone dose calculated	
Nurse in charge aware Name:	
Consultant in charge aware Name:	

Documentation of any adverse events: