Tetracosactide (Tetracosactrin)

****This monograph is under review****

Neonatal

FORM

Ampoule containing 250microgram in 1ml

DO NOT USE THE DEPOT INJECTION.

INDICATION

Used as a diagnostic agent for adrenal function.

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth – 6 months	36 micrograms/kg (Maximum dose 250 micrograms)	Single dose at time zero	IV

Special care is needed when calculating the dose.

RECONSTITUTION

Already in solution

DILUTION

For all low dose Synacthen ® tests, dilution of the injection will be required.

Tetracosactide 250 microgram/ml inj.	1ml
Sodium Chloride 0.9%	Up to 10ml total

This gives a 25microgram in 1ml solution. Use the required volume

METHOD OF ADMINISTRATION

IV injection

COMPATIBILITY

No information

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

- See Summary of Product Characteristics and most recent edition of BNF for Children

SPECIAL MONITORING REQUIREMENTS

Consult current edition of Dynamic Function Test Handbook for Clinicians(SPEG MCN)/ liaise with Endocrine for sampling times/interpretation of results

To determine which tests are to be performed and which resulting samples are to be obtained consult with Endocrinology Consultant at RHC before initiating tests.

FURTHER INFORMATION

LICENSED STATUS

Licensed status: Licensed for use as a diagnostic test for adrenal function. Doses used in this monograph may not be licensed.

APPLICABLE POLICIES

West of Scotland Neonatal Guidelines:

Managed Clinical Network of Scottish Paediatric Endocrine Group(SPEG MCN) Dynamic Function Tests for use in Paediatric Endocrinology V2 1.9.2019:

https://www.speg.scot.nhs.uk/wp-content/uploads/2020/03/9.-SPEG-Dynamic-Function-Tests-in-Paediatric-Endocrinology-2018GS-Draft-3-1.pdf

Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared by:	WoS Neo pharm group	Checked by	Maria Tracey
Date prepared	May 2016	Date updated	May 2020
Updated by	Maria Tracey	Review Date	May 2023
Extension Agreed By	WoS NPG	Review Date	May 2027

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Network. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department