Levofloxacin

****This monograph is under review****

Neonatal

FORM Vial containing 500mg in 100ml

INDICATION Treatment of susceptible organism on microbiology advice

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth – 6 months	8mg/kg/dose	Every 12 hours	IV

RECONSTITUTION Already in solution.

DILUTION Dilution not required

Dosage guide

- 5mg in 1ml
- 8mg in 1.6ml

METHOD OF ADMINISTRATION IV infusion over 30 minutes

COMPATIBILITY

Solution compatibility	Sodium chloride 0.45%, Sodium chloride 0.9% , Glucose 5%, Glucose 10%, Parenteral Nutrition
Solution incompatibility	
IV Line compatibility	
IV Line incompatibility	

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

SPECIAL MONITORING REQUIREMENTS

FURTHER INFORMATION

PH 3.8 – 5.8

LICENSED STATUS Unlicensed in neonates

NEONATAL West of Scotland Drug Monographs
Parenteral Route

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	001	Supersedes:	Nil
Prepared by:	Peter Mulholland	Checked by	Wos Neonatal Pharmacists
Date prepared	April 2018	Date updated	
Updated by		Review Date	April 2021
Extension Agreed By	WoS NPG	Review Date	July 2027

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.
Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the package insert, BNF for Children, and Summary of Product Characteristics. For further advice contact your clinical pharmacist or pharmacy department.