FLUCLOXACILLIN

FORM Vials containing 250mg, 500mg, 1g powder for reconstitution

INDICATION a) Treatment of infection where sensitivities indicate. Suspected staphylococcal infections

b) Osteomyelitis, cerebral abscess, staphylococcal meningitis

DOSE RANGE

a) Treatment of infection where sensitivities indicate. Suspected staphylococcal infections

AGE	DOSE	FREQUENCY	ROUTE
< 7 days	25mg/kg*	12 hourly	I.V.
7 - 20 days	25mg/kg*	8 hourly	I.V.
21 -28 days	25mg/kg*	6 hourly	I.V.
1-6 months	12.5-25mg/kg*	6 hourly	I.V.

*Doses should be doubled in severe infection

b) Osteomyelitis, cerebral abscess, staphylococcal meningitis

AGE	DOSE	FREQUENCY	ROUTE
< 7 days	50-100mg/kg	12 hourly	I.V.
7 - 20 days	50-100mg/kg	8 hourly	I.V.
21 -28 days	50-100mg/kg	6 hourly	I.V.
1-6 months	50mg/kg	6 hourly	I.V.

RECONSTITUTION

250mg vial: (average) displacement value 0.2ml.
Reconstitute with 4.8ml water for injection

500mg vial: (average) displacement value 0.4ml.
Reconstitute with 9.6ml water for injection

1g vial: (average) displacement value 0.7ml.
Reconstitute with 19.3ml water for injection

Resulting Solution for all vials 50mg/ml

DILUTION Not required

METHOD OF ADMINISTRATION Administer required volume of the reconstituted solution over 3-5 minutes

Neonatal

COMPATIBILITY

Solution compatibility	Glucose 5%, Sodium chloride 0.9%
Solution incompatibility	No Information
IV Line compatibility	No Information
IV Line incompatibility	Amikacin, Amiodarone, Atropine, Calcium Gluconate, Ciprofloxacin, Clarithromycin, Dobutamine, Erythromycin, Gentamicin, Midazolam, Morphine, Lipid emulsions

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION Each 1g of powder contains 2 - 2.26mmol of Sodium.

PH 5-7

LICENSED STATUS Licensed

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared By/Checked By	Jacqueline Rooney / Susan Kafka	Final Approval By	WoS Neonatal Pharmacist Group
Date Prepared	August 2026	Review Date	August 2029

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department