Carnitine

****This monograph is under review****

FORM Paediatric Oral Solution 300mg/ml
** Special Care required – other strengths available**

INDICATION Treatment of Primary and Secondary Carnitine Deficiency

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth to 6 months	Initially 25mg/kg/dose*	FOUR times daily	Oral
	May increase to 175mg/kg/day or 3g daily		

* Specialist consultants may recommend different doses on an individual baby basis.

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

- L-Carnitine is an amino acid derivative which is an essential co-factor of fatty acid metabolism.
- Dose over 100mg/kg/day may cause diarrhoea.
- Can be diluted in water if required.

LICENSED STATUS Licensed in neonate and infants under 12 years

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	001	Supersedes:	Nil
Prepared by:	WoS neo pharm group	Checked by	Maria Tracey
Date prepared	Aug 2016	Date updated	
Updated by		Review Date	Aug 2019
Extension Agreed By	WoS NPG	Review Date	August 2027

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist network. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.