# Fluconazole

**This monograph is under review**

**FORM** Vial containing 50mg in 25ml
Infusion bag containing 100mg in 50ml

**INDICATION** Treatment of deep systemic fungal infections
Severe Life Threatening Infections, Meningitis
Prophylaxis of fungal infections

**DOSE RANGE**

| AGE | DOSE | FREQUENCY | ROUTE |
|---|---|---|---|
| < 2 weeks | 12mg/kg | Every 72 hours | IV |
| 2 - 4 weeks | 12mg/kg | Every 48 hours | IV |
| > 4 weeks | 12mg/kg | Once daily | IV |
| * In renal impairment - reduce dose to 6mg/kg if Creatinine > 80 | | | |

**RECONSTITUTION** Already in solution

**DILUTION** If further dilution is necessary, dilute with Sodium Chloride 0.9% or Glucose 5%.

**METHOD OF ADMINISTRATION** Infuse required volume over 15 minutes

**COMPATIBILITY**

| | |
|---|---|
| **Solution compatibility** | Glucose 5%, Glucose 10%, Sodium chloride 0.45%, Sodium chloride 0.9% |
| **Solution incompatibility** | No information |
| **IV Line compatibility** | Aciclovir, amikcain, aminophylline, amiodarone, benzylpenicillin, dexamethasone, dobutamine, dopamine, gentamicin, heparin sodium, meropenem, midazolam, morphine, piperacillin with tazobactam, ranitidine, tacrolimus, tazocin, tobramycin vancomycin, vecuronium zidovudine.<br>Compatible with the following medicines when they are ready-diluted: gangciclovir, linezolid, metronidazole. |
| **IV Line incompatibility** | Amphotericin, calcium gluconate, cephalosporin antibiotics, clindamycin,<br>co-trimoxazole, diazepam, digoxin, erythromycin, furosemide, imipenem. |

**THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED**

**CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS**
- See Summary of Product Characteristics and most recent edition of BNF for Children

**FURTHER INFORMATION** Fluconazole may potentiate the effects of Warfarin and also increase blood levels of Phenytoin, Theophylline and Ciclosporin.

**PH** 4 - 8

**LICENSED STATUS** Licensed

**APPLICABLE POLICIES** West of Scotland Neonatal Guidelines

Consult local policy if applicable

# NEONATAL West of Scotland Drug Monographs
## Parenteral Route

| | | | |
|---|---|---|---|
| Document Number: | 001 | Supersedes: | Nil |
| Prepared by: | WoS Neo pharm group | Checked by | Michael Mitchell |
| Date prepared | May 2016 | Date updated | September 2019 |
| Updated by | Lauren Williams | Review Date | September 2022 |
| Extension Agreed By | WoS NPG | Review Date | April 2027 |

**Administer reconstituted solutions immediately.**
**All vials, ampoules and infusion bags are for single use only unless otherwise stated.**

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Network. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department