Levetiracetam

FORM Oral solution containing 100mg/ml

INDICATION Treatment of neonatal seizures

STARTING MAINTENANCE DOSE

GESTATIONALAGE	AGE	DOSE	FREQUENCY	ROUTE
<36 WEEKS	< 7 days	5-10mg/kg*	Every 8 hours	Oral
	≥ 7 days	10-20mg/kg*	Every 12 hours	
≥36 WEEKS	< 7 days	7.5-10mg/kg*	Every 8 hours	Oral
	≥7 days	10-30mg/kg*	Every 12 hours	

Startingdoses/standard dosing ranges only. Doses up to 60mg/kg/day (pre-term infants) and 100mg/kg/day (term infants) have been given in some studies (on Specialist Neurology advice only)

CAUTION in renal impairment – maintenance dose adjustments required. Discuss with Pharmacy

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION Excipient contents vary between products. The use of the Milpharm brand should be avoided where possible, however the benefit of treatment may still out-weigh risk

STORAGE Room temperature

LICENSED STATUS Licensed from ONE month of age

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared/Checked by:	Susan Kafka/Frances Lowrie/Marie Stewart	Final approval by	WoS Neonatal Pharmacist Group
Date prepared	September 2026	Review date	September 2029

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.