Vitamin D (Colecalciferol)

FORM

a) Oral drops containing 2400units per 1ml
b) Oral drops containing 400units per drop

INDICATION

1) For Vitamin D deficiency
2) Routine supplementation in babies > 34 weeks GA who are breastfeeding or on < 500ml a day of formula milk.

DOSE RANGE

1) For Vitamin D deficiency as per Metabolic Bone Disease guideline

INDICATION	DOSE	FREQUENCY	ROUTE
PTH high (>7 pmol/L) and vitamin D 25-50nmol/l	400units	ONCE daily for 6 weeks	Oral
PTH high (>7 pmol/L) and vitamin D < 25nmol/l	2400units*	ONCE daily for 12 weeks	Oral
PTH normal (2-7 pmol/L) and vitamin D < 50nmol/l	400units	ONCE daily for 6 weeks	Oral

- **Above dosing is in line with the metabolic bone disease guideline and assumes appropriate calcium or phosphate supplementation.**
- **The above dose is in addition to any Vitamin D the baby may be receiving with standard pre-term supplements**
- ***It may be appropriate to consider doses of 1200units in certain clinical situations**

2) Routine supplementation in babies > 34 weeks GA who are breastfeeding of on < 500ml a day of formula milk (some centres use Pro Health Drops – see below)

AGE	DOSE	FREQUENCY	ROUTE
>34 weeks Breast fed / <500ml formula	400units	Once daily	Oral

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

- Administer dose with feeds where possible
- Dropper should be removed from product prior to discharge
- In renal impairment use hydroxylated equivalent (alfacalcidol)
- Pro Health Drops are the recognised food supplement recommended by the Scottish Government for vitamin D supplementation. 400 units is equivalent to 10micrograms and is administered in one drop of the solution.

NEONATAL West of Scotland Drug Monographs
ORAL/OTHER ROUTES

LICENSED STATUS

Document Number:	001	Supersedes:	Colecalciferol_WOSOOR
Prepared by/Checked By	Lauren Williams/Frances Lowrie/Jacqueline Rooney	Final Approval By:	WoS Neonatal Pharmacist Group
Date prepared	September 2026	Review Date	September 2029

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.