Naloxone Intravenous

FORM Ampoule containing 400 micrograms/ml injection

INDICATION

1. Overdose with opioids
2. Reversal of respiratory and CNS depression in neonate following maternal opioid use during labour

1 Overdose with opioids

AGE	DOSE	FREQUENCY	ROUTE
Birth to 6 months	Initially 1 – 10 micrograms/kg, if no response repeat at intervals of 1 minute up to 5 times. If still no response give a single dose of 100micrograms/kg then review diagnosis if still not responding	As required	IV

2 Reversal of respiratory and CNS depression in neonate following maternal opioid use during labour

AGE	DOSE	FREQUENCY	ROUTE
Birth to 6 months	10micrograms/kg Repeat every 2-3minutes if required Further doses may be required if respiratory function deteriorates following initial response	As required	IV

In neonates with respiratory depression by maternal opioids, it is generally preferred to give an IM injection – see Naloxone IM monograph

RECONSTITUTION Already in solution

DILUTION

Take 0.5ml of the 400microgram/ml Naloxone Injection.
Dilute with 0.9% Sodium Chloride injection to a total of 10ml

This gives a 20 microgram/ml solution.

Dosage guide for diluted solution

- 10 micrograms in 0.5ml
- The starting dose of 10 microgram/kg = 0.5 ml/kg of above solution

METHOD OF ADMINISTRATION Rapid IV injection

COMPATIBILITY

Solution compatibility	Glucose 5%, sodium chloride 0.9%
Solution incompatibility	All other solutions
IV Line compatibility	No other drugs or fluids in same line at same time
IV Line incompatibility	All other drugs

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION Naloxone given IV should produce an effect within 2-3minutes and should last 3-4hours.

Caution – the half-life of a narcotic drug may be longer than that of naloxone.

PH pH ranges from 3 to 4.5

LICENSED STATUS Licensed

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Version	002	Supercedes	001
Prepared/checked by	Olivia Scales/Susan Kafka/Natalie Donald	Approved by	WoS Neonatal Pharmacist Group
Updated	September 2026	Review Date	September 2029

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department