Amoxicillin

FORM Vial containing 250mg powder for reconstitution.

INDICATION

1. Penicillin antibiotic used in the treatment of infection where sensitivities or guidelines dictate.

2. Severe infections including Listerial meningitis, group B streptococcal infection and enterococcal endocarditis.

3. Babies re-admitted from home, treated for community acquired neonatal sepsis, in combination with cefotaxime and gentamicin.

DOSE RANGE

1. Penicillin antibiotic used in the treatment of infection where sensitivities or guidelines dictate ; management of uncomplicated group B streptococcal infection

AGE	DOSE	FREQUENCY	ROUTE
< 7days	50mg/kg per dose	every 12 hours	IV
7days – 6 months	50mg/kg per dose	every 8 hours	IV

*dose may be escalated as below if concerns over failure to respond

2. Severe infections including meningitis caused by Listeria or group B streptococcal infection; and in enterococcal endocarditis (in combination with another antibiotic)

AGE	DOSE	FREQUENCY	ROUTE
< 7days	100mg/kg per dose	every twelve hours	IV
7 – 28 days	100mg/kg per dose	every eight hours	IV
1 – 6 months	50mg/kg per dose	every 4 hours	IV

3. Babies re-admitted from home, treated for community acquired neonatal sepsis, in combination with cefotaxime and gentamicin.

AGE	DOSE	FREQUENCY	ROUTE
< 7days	100mg/kg per dose	every twelve hours	IV
7 – 28 days	100mg/kg per dose	every eight hours	IV
1 – 6 months	50mg/kg per dose	every 4 hours	IV

For positive cultures where the isolate is reported as 'I' an increased exposure dosing regimen is required. Exposure may be increased as a function of dose, frequency or administration time depending on the drug.
Please refer to the following guidance *"Isolates reported as 'I' (susceptible increased exposure)- dosing schedule for neonates and children"* or contact Pharmacy for further information.

RECONSTITUTION Add 2.3ml water for injection BP to each 250mg vial and shake gently until completely dissolved, to produce a 100mg/ml solution.

DILUTION No further dilution required.

Neonatal

NEONATAL West of Scotland Drug Monographs
Parenteral Route

METHOD OF ADMINISTRATION	**Intermittent IV Infusion** *(dose >30mg/kg associated with CNS toxicity)* Further dilute with sodium chloride 0.9% or glucose 5% to convenient volume and give over 30minutes Note: administration of doses 30-50mg/kg by slow IV bolus is out with the SmPC however has historically been done in neonatal practice. This may be required for administration on post natal wards **IV bolus over 3-5 minutes** (see note below) No further dilution required. Give as a slow bolus

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

PH 8.0-10.0

LICENSED STATUS Licensed.

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared by/Checked by:	Susan Kafka/Andrew Stockton/Roisin McCabe	Approved by	WoS Neonatal Pharmacist Group
Date prepared	June 2026	Review date	June 2029

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications. This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement. For further advice please contact your clinical pharmacist or pharmacy department