Spironolactone

FORM Oral solution 50mg/5ml

INDICATION Prevention of hypokalaemia whilst on diuretic therapy
Oedema in heart failure and in ascites

DOSE RANGE

AGE	DOSE*	FREQUENCY	ROUTE
Neonate	500micrograms-1mg/kg	12 hourly	Oral
1month- 6months	500micrograms-1.5mg/kg	12 hourly	Oral

*Higher doses of up to 7mg/kg/day in neonates or up to 9mg/kg/day in children 1 month and over may be used in resistant ascites (discuss with consultant before use)

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION Caution required in renal impairment or if patient is also prescribed potassium supplements or drugs likely to raise potassium levels. Monitor potassium levels closely.

LICENSED STATUS Unlicensed special

APPLICABLE POLICIES West of Scotland Neonatal Guidelines
Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared by/Checked by:	Jacqueline Rooney/Marie Stewart/Frances Lowrie	Final approval by:	WoS Neonatal Pharmacist Group
Date prepared	September 2026	Review date:	September 2029

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.