Paracetamol – Analgesia/Pyrexia

See most recent edition on BNFC for advice on maximum doses and treatment of therapeutic excess

FORM Oral Suspension 120mg/5ml (caution other strengths available)

INDICATION

1. Pain Relief, Pyrexia
2. **Prophylaxis of post-immunisation pyrexia following immunisation with meningococcal group B vaccine (Bexsero®)**

DOSE RANGE

1. **Pain relief, pyrexia**

AGE	DOSE	FREQUENCY	ROUTE
Neonate 28-32 weeks gestation	10-15mg/kg (Max 30mg/kg in 24 hours)	Every 8-12 hours	Oral
Neonate > 32 weeks gestation	10-15mg/kg (Max 60mg/kg in 24 hours)	Every 6-8 hours	
Child 1 month (or 44 weeks CGA) - 6 months	10-15mg/kg (Max 60mg per dose)	Every 6-8 hours (Max 4 doses in 24 hours)	

2. **Prophylaxis of post-immunisation pyrexia following immunisation with meningococcal group B vaccine (Bexsero®)**

AGE	DOSE	FREQUENCY	ROUTE
Neonate 28-32 weeks gestation	10-15mg/kg (Max 30mg/kg in 24 hours)	Every 8 hours*	Oral
Neonate > 32 weeks gestation	10-15mg/kg (Max 60mg per dose and Max 60mg/kg in 24 hours)	Every 6 hours*	
Child 1 month (or 44 weeks CGA) - 6 months	10-15mg/kg (Max 60mg per dose)	Every 6 hours*	

* For 3 doses, first dose to be given at the time of or as soon as possible after vaccination

Please refer to the BNFc for alternative routes if IV or oral dosing is not suitable

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

- Use with caution in liver dysfunction or where there are risk factors for hepatotoxicity
- Do not give with any other preparation containing paracetamol
- Caution in unconjugated hyperbilirubinemia due to risk of accumulation. Consider reducing dose and/or increasing the dosage interval

Neonatal

NEONATAL West of Scotland Drug Monographs
ORAL/OTHER ROUTES

LICENSED STATUS	Oral suspension not licensed for use in children under 2 months
APPLICABLE POLICIES	West of Scotland Neonatal Guidelines Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared By/Checked By	Julie Lowther/Frances Lowrie/Lauren Williams/Olivia Scales	Final Approval By	WoS Neonatal Pharmacist Group
Date Prepared	September 2026	Review Date	March 2028

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.