Sugammadex

FORM Vial containing sugammadex sodium 200 mg in 2 ml

INDICATION Reversal of neuromuscular blockade

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Neonate – 6 months	2 mg/kg	Single dose Repeated under consultant advice according to response	IV

Neonatal

RECONSTITUTION Already in solution

DILUTION

Sugammadex 200 mg in 2 ml	1 ml
Sodium chloride 0.9%	Up to 10 ml

Gives a 10 mg/ml solution. Use the required volume.

METHOD OF ADMINISTRATION Rapid IV bolus injection

Rapidly flush the line with sodium chloride 0.9%

COMPATIBILITY

Solution compatibility	Sodium chloride 0.45%, sodium chloride 0.9%, glucose 5%
Solution incompatibility	No information
IV Line compatibility	Do not bolus via a line containing any other medicines
IV Line incompatibility	No information

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS See Summary of Product Characteristics and BNF for Children

FURTHER INFORMATION Sugammadex has a half-life of 2 hours therefore if rocuronium or vecuronium needs re-initiated with 24 hours dose titration may be needed.

LICENSED STATUS Unlicensed in neonates

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared By/Checked By	Andrew Stockton/Lauren Williams/Natalie Donald	Final Approval By	WoS NPG
Date Prepared	September 2026	Review Date	September 2029

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department