Sodium Acid Phosphate

FORM Oral solution containing phosphate 1mmol/ml and sodium 1mmol/ml

INDICATION Prevention of Metabolic Bone Disease in Premature neonates < 32 weeks
.

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth to 6 months *When PTH is within normal limits (2-7 pmol/L)*	0.5mmol/kg/dose Adjust dose according to response up to maximum of 1.5mmol/kg/dose)	TWICE daily	Oral

- **Above dosing is in line with the Metabolic Bone Disease guideline and assumes appropriate calcium supplementation has been prescribed +/- appropriate vitamin D supplementation as per guideline**

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

- May also started where biochemistry shows hypophosphataemia after discussion with neonatal consultant
- Use with caution in renal failure
- Treatment will usually stop prior to discharge
- Available preparations may contain excipients which are not suitable for neonates, such as ethanol or chloroform. Please check suitability of product before use.
- Do not administer at the same time as calcium supplements as this can cause poor absorption. Administer separately and between feeds where possible.

STORAGE Room temperature. Store in fridge once opened. Some products have a seven day expiry once opened – please check bottle for guidance.

LICENSED STATUS Unlicensed special product

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared By/Checked By	Lauren Williams/Marie Stewart/Andrew Stockton	Final Approval By	WoS Neonatal Pharmacist Group
Date Prepared	September 2026	Review Date	September 2029

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.