Metronidazole

****This monograph is under review****

FORM Oral Suspension 200mg/5ml

INDICATION Treatment of Necrotising Enterocolitis (when oral absorption considered normal)
Treatment of other infections where guidelines or sensitivities indicate

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth to 2 months	7.5mg/kg	TWICE daily	Oral
2months – 6 months	7.5mg/kg	THREE times daily	Oral

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION Contains metronidazole benzoate
Contains sorbitol

LICENSED STATUS Licensed in all ages

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:
Consult local policy if applicable

Neonatal

Document Number:	002	Supersedes:	001
Prepared by:	WoS Neo pharm group	Checked by	June Grant
Date prepared	Aug 2016	Date updated	August 2021
Updated by	Roisin McCabe	Review Date	August 2024
Extension Agreed By	WoS NPG	Review Date	July 2027

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist network. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.