Clonidine

FORM Ampoule containing 150 micrograms in 1 ml

INDICATION Sedation, analgesia, opioid withdrawal

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth – 6 months	0.5-1 micrograms/kg/h (Usual max dose 2 micrograms/kg/h)	Continuous infusion	IV

PRESCRIPTION OF CONTINUOUS INFUSION

200 micrograms/kg in 50ml infusion fluid
This gives:

- 0.5 micrograms/kg/h at 0.13 ml/h
- 1 microgram/kg/h at 0.25 ml/h

RECONSTITUTION

Already in solution

DILUTION

Clonidine 150 micrograms/ml	1.33 ml/kg*
Sodium chloride 0.9%	Up to 50 ml

*1.33 m/kg equivalent to 200 micrograms/kg

METHOD OF ADMINISTRATION

For continuous Infusion
By continuous intravenous infusion, flow rate adjusted according to the baby's response (see prescription section for details).

Clonidine must not be bolused

COMPATIBILITY

Solution compatibility	Sodium chloride 0.9%, glucose 5%, glucose 10%
Solution incompatibility	No information
IV Line compatibility	Adrenaline, dobutamine, dopamine, fentanyl, furosemide, insulin, heparin, ketamine, isoprenaline, labetalol, midazolam, milrinone, morphine, noradrenaline, potassium chloride, sildenafil, sodium nitroprusside (in glucose only).
IV Line incompatibility	Omeprazole

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION	Monitor heart rate and blood pressure
LICENSED STATUS	Unlicensed use in neonates
APPLICABLE POLICIES	West of Scotland Neonatal Guidelines Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared By/Checked By:	Andrew Stockton/Lauren Williams/Susan Kafa	Final Approval By:	WoS Neonatal Pharmacist Group
Date Prepared:	September 2026	Review Date:	September 2029

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department