Magnesium Aspartate (Magnaspartate)

FORM Magnaspartate contains 243mg/10mmol Mg^{2+} per sachet, powder for oral solution

INDICATION Treatment and prevention of magnesium deficiency

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
0 – 6 months	0.2mmol/kg	THREE times daily	Oral

RECONSTITUTION

Magnaspartate	1 sachet
Water for Injection	10mls*

- Gives Mg^{2+} 1mmol in 1ml solution. Use the required volume

*10ml volume for dilution is off-label but established clinical practice due to larger volumes not being well tolerated in this age group

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION Monitor serum magnesium throughout treatment

Discard any remaining solution after dose is administered.

STORAGE Store at room temperature

LICENSED STATUS Off-label use of licensed medicine

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	001	Supersedes:	Nil
Prepared By/Checked By	Olivia Scales/Frances Lowrie/Marie Stewart	Final Approval By	WoS Neonatal Pharmacist Group
Date Prepared	September 2026	Review Date	September 2029

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.