Caffeine Citrate

NOTE : Take special care with calculating doses and administration volumes. A ten times overdose could prove fatal.

FORM

Ampoule containing 20mg/ml caffeine citrate solution for oral administration

PRESCRIBE AS CAFFEINE CITRATE

INDICATION

1. Treatment and prophylaxis of apnoea of prematurity
2. Prevention of bronchopulmonary dysplasia

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Neonate - 6 months	20mg/kg	**STAT LOADING DOSE**	Oral
Neonate - 6 months	10mg/kg	Once daily **MAINTENANCE***	Oral

- *Commence maintenance 24 hours after loading dose
- Consider 10mg/kg twice daily dosing when on concomitant enzyme inducing drug e.g. phenobarbital
- Infants over 28 days who are still apnoeic should benefit from maintenance caffeine doses administered twice daily
- It may be appropriate to extend caffeine treatment duration for certain infants. Please consult local policy and guidelines

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

See Summary of Product Characteristics and most recent edition of BNF for Children

Review dose if tachycardia occurs.

FURTHER INFORMATION

- Oral doses should be drawn up using appropriate filter straws
- Store at room temperature
- Caution in impaired renal or hepatic function
- See summary of product characteristics for signs of toxicity

LICENSED STATUS

Off-label use of licensed product

NEONATAL West of Scotland Drug Monographs
ORAL/OTHER ROUTES

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	004	Supersedes:	003
Prepared by:	Lauren Williams/Andrew Stockton/ Frances Lowrie	Final Approval By:	WoS Neonatal Pharmacist Group
Date Prepared	September 2026	Review Date	September 2029

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.