Albumin

FORM Vials containing 20% albumin

INDICATION Consider in symptomatic hypoalbuminaemia (Albumin <20g/L)

DOSE

Symptomatic hypoalbuminaemia

AGE	DOSE	FREQUENCY	ROUTE
0 – 6 months	5ml/kg (equal to 1g/kg)	Once daily	IV infusion

RECONSTITUTION Already in solution

DILUTION Further dilution not required

METHOD OF ADMINISTRATION Intravenous infusion over 60minutes

Can be used as a continuous infusion in some circumstances

Neonatal

COMPATIBILITY

Solution compatibility	Sodium chloride 0.9%, Glucose 5%, Glucose 10%
Solution incompatibility	Water for injection
IV Line compatibility	Do not administer with other drugs
IV Line incompatibility	Do not administer with other drugs

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

SODIUM CONTENT Sodium Content variable – average 0.12-0.16mmol/ml

LICENSED STATUS Licensed

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared/Checked by:	Susan Kafka/ J Rooney/Frances Lowrie	Approved by	WoS Neonatal Pharmacist Group
Date updated	September 2026	Review Date	September 2029

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department