Adrenaline (Epinephrine)- for RESUSCITATION 1:10,000

MORE THAN ONE STRENGTH AVAILABLE – ENSURE THE CORRECT STRENGTH IS USED FOR THE APPROPRIATE INDICATION

Neonatal

BRAND NAME Adrenaline (Epinephrine)

FORM Ampoule/pre-filled syringe containing 100microgram/ml (**1 in 10,000**)
CAUTION – ENSURE CORRECT STRENGTH SELECTED

INDICATION Acute Resuscitation

Acute Resuscitation
DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth - 6 months	20 microgram/kg (0.2ml/kg of **1:10,000**)	1st dose	I.V. rapid bolus / intraosseous / UVC* See below for Endotracheal(ET) administration
	Repeat above dose every 3-5minutes if required and consider use of sodium bicarbonate		

***The UK Resuscitation Council advises administration of adrenaline by umbilical venous catheter as route of choice in resuscitation of the newborn.**

ET administration

Doses of adrenaline only may be given by Endotracheal (ET) tube, if no venous access. For ET tube administration inject quickly down a narrow-bore suction catheter beyond the tracheal end of the ET tube and then flush with 1 or 2 ml Sodium Chloride 0.9% injection. If the tracheal route is used, a dose of 100microgram/kg (1ml/kg of **1:10,000**) should be given

RECONSTITUTION Already in solution

DILUTION Dilution not required

METHOD OF ADMINISTRATION By rapid intravenous bolus / intraosseous / UVC administration.
If given intraosseous, flush with 1 or 2 ml Sodium Chloride 0.9% injection.
See above for ET administration

COMPATIBILITY

Solution compatibility	Sodium chloride 0.9%, Glucose 5%, Glucose 10%,
Solution incompatibility	No information
IV Line compatibility	Atracurium, Calcium Gluconate, Caffeine citrate, Dexmedetomidine, Dobutamine, Dopamine, Fentanyl, Glyceryl Trinitrate, Heparin, Midazolam, Milrinone, Morphine, Noradrenaline, Potassium Chloride, Rocuronium, Vecuronium Compatible in Sodium Chloride 0.9% only - Insulin (soluble)
IV Line incompatibility	Aminophylline, Epoprostenol, Omeprazole, Sodium bicarbonate

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

Monitoring requirements:
Heart rate and rhythm, arterial blood pressure and peripheral perfusion.
Urine output and renal function.

STORAGE

Do not use discoloured solutions or solution containing a precipitate.

PH

2.2 – 4.0

LICENSED STATUS

Some preparations may be unlicensed

APPLICABLE POLICIES

West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared by/Checked by:	Susan Kafka/ Jacqueline Rooney/Natalie Donald	Approved by	WoS Neonatal Pharmacist Group
Date prepared	June 2026	Review Date	June 2029

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department