Sodium Chloride

FORM Oral Solution 29.22% (containing 5mmol/ml Sodium)

INDICATION Treatment of hyponatraemia

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth to 6 months	As per calculation below. Adjust according to response OR 1 mmol/kg/dose	Give daily requirement in divided doses* THREE times daily	Oral

* It is preferable to give smaller doses more often (in milk feeds) as this is often better tolerated by babies.

Sodium deficit (mmol) = (135 – plasma sodium) x 0.6 x body weight (kg)

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION It is usual practice to round sodium doses to the nearest 0.5ml for ease of administration.

STORAGE Storage conditions and shelf life vary depending on product and if opened or sealed. Consult bottle for details.

LICENSED STATUS Licensed and unlicensed 'special' preparations available.

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared by /Checked by	Andrew Stockton/Julie Lowther/Marie Stewart	Final approval by:	WoS Neonatal Pharmacy Group
Date updated	September 2026	Review Date	September 2029

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.