Valganciclovir

Handle as a cytotoxic as considered a potential teratogen and carcinogen in humans
Use aprons and gloves for preparation and administration

FORM 250mg in 5ml oral solution (dry powder for reconstitution)

INDICATION Treatment of congenital CMV

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Neonate >32weeks gestation to 6months*	16mg/kg	12 hourly	Oral

*Dosing for clinical indications for babies less than 32 weeks gestation should be discussed with pharmacy

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

- Follow directions on side of bottle for reconstitution
- Pregnant staff should not handle this product.
- If powder or reconstituted solution come into contact with skin or mucosa, wash off immediately with soap and water. Avoid inhalation of the powder.
- Oral syringes and nappies should be discarded as cytotoxic waste
- Use with caution in renal impairment. Dose reduction may be needed. Contact pharmacy for further information.
- Can cause neutropenia, thrombocytopenia and anaemia. Monitor blood and platelet counts.
- Routine therapeutic drug monitoring is not required however may be considered if there are suspicions of toxicity.

STORAGE Once reconstituted the solution should be refrigerated. Use within 49 days of reconstitution

LICENSED STATUS Unlicensed use.

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared by/Checked By	Roisin McCabe/J Rooney/F Lowrie	Final Approval By	WoS Neonatal Pharmacist Group
Date Prepared	September 2026	Review Date	September 2029

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.