Ibuprofen

FORM Ampoule containing 10mg in 2ml

INDICATION Treatment of a haemodynamically significant patent ductus arteriosus in preterm newborn infants less than 34 weeks of gestational age.

DOSE RANGE **3 dose course (including subsequent courses)**

AGE	DOSE	FREQUENCY	ROUTE
Preterm neonate < 34 weeks gestation	10mg/kg	LOADING DOSE (give after the first 6hr of life) 1 dose only followed at 24 hour intervals by	IV
	5mg/kg	24hours after loading dose 2nd and 3rd doses given at 24 hour intervals*	IV

* If the ductus arteriosus does not close 48 hours after the last injection or if it reopens, a second course of 3 doses, as above, may be given. If the condition is unchanged after the second course of therapy, surgery of the patent ductus arteriosus may then be necessary.

Alternatively, if duct remains open 24 hours after the third dose, a fourth and fifth dose given at 24 hour intervals may be considered under the direction of a consultant.

Consider oral dosing for babies tolerating >100ml/kg/day enteral feed

RECONSTITUTION Already in solution

DILUTION Preferably to be administered undiluted

METHOD OF ADMINISTRATION Administer as a short IV infusion over 15 minutes.

Before and after administration, to avoid contact with any acidic solution, flush infusion over 15 minutes with 1.5 to 2ml of either Sodium chloride 0.9% or Glucose 5% solution for injection.

COMPATIBILITY

Solution compatibility	sodium chloride 0.45%, sodium chloride 0.9%, glucose 5%
Solution incompatibility	All other IV fluids.
IV Line compatibility	No other drugs or IV fluids at same time in same line.
IV Line incompatibility	All other drugs

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

- Avoid concomitant use with steroids due to increased risk of spontaneous GI perforation. If concomitant use is unavoidable this should be under the direction of a Consultant only.
- Extravasation risk- may cause irritation to surrounding tissues upon extravasation.
- Chlorhexidine must not be used to disinfect the neck of the amp as it is not compatible with Pedea ® For asepsis use ethanol 60% or isopropyl alcohol 70%.
- Use with caution in renal, cardiac or hepatic impairment. Careful monitoring of renal and gastrointestinal function is recommended during treatment
- Monitor for signs of bleeding

PH 7.8 – 8.2

LICENSED STATUS Licensed medicine

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	004	Supersedes:	003
Updated by/Checked by:	Roisin McCabe/Olivia Scales/Susan Kafka	Approved by:	WoS Neonatal Pharmacist Group
Date updated	August 2026	Review date	February 2028

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department