Omeprazole

This monograph is under review

Neonatal

FORM 40mg vial – powder for reconstitution

INDICATION Gastro-oesophageal reflux disease

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth – 6 months	Initially 500micrograms/kg. Increased to 2mg/kg once daily if necessary (max 20mg).	Once daily	IV infusion

RECONSTITUTION

1. Reconstitute the 40mg vial with 5ml sodium chloride 0.9% to give an 8mg/ml solution.

2. Mix thoroughly making sure all omeprazole has dissolved.

DILUTION

Omeprazole 8mg/ml solution	1ml
Sodium chloride 0.9%	Up to 20ml

Gives a 0.4mg in 1ml solution. Use the required volume.

METHOD OF ADMINISTRATION Administer the required volume of diluted solution over 20 - 30 minutes.

COMPATIBILITY

Solution compatibility	Sodium chloride 0.9%, Glucose 5%,
Solution incompatibility	No info
IV Line compatibility	No info
IV Line incompatibility	Very restricted: including (but not limited to) adrenaline, amiodarone, atracurium, clonidine, dobutamine, dopamine, esmolol, fentanyl, furosemide, heparin, insulin, labetalol, midazolam, morphine, noradrenaline, rocuronium, sodium nitroprusside, vancomycin, vecuronium.

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION Do not use if any particles are present in the solution.

STORAGE	Powder: do not store above 25℃. Keep vials in the outer carton in order to protect from light. Reconstituted solution: Use immediately
PH	8.5 – 10.5 (once reconstituted)
LICENSED STATUS	Unlicensed use
APPLICABLE POLICIES	West of Scotland Neonatal Guidelines: Consult local policy if applicable

Document Number:	004	Supersedes:	003
Prepared by:	Andrew Stockton	Checked by	WOS Neo Pharmacy Group
Date Prepared	November 2023	Review Date	November 2026
Extension Agreed By	WoS NPG	Review Date	November 2027

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Network. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department