Salbutamol

FORM Ampoule containing 5 mg in 5 ml

INDICATION Severe hyperkalaemia

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Neonate – 6 months	4 micrograms/kg	Single Dose	IV

Dose may be repeated according to response

RECONSTITUTION Already in concentrated solution

DILUTION

Salbutamol 5 mg in 5 ml solution	0.2 ml
Sodium chloride 0.9%	Up to 20 ml
This gives 200 micrograms in 20 ml (10 micrograms/ml)	

METHOD OF ADMINISTRATION Slow IV bolus over 5 minutes.
4 micrograms/kg = 0.4 ml/kg of diluted solution.

COMPATIBILITY

Solution compatibility	Sodium chloride 0.9%, Glucose 5%
Solution incompatibility	No information
IV Line compatibility	Ideally, salbutamol bolus should be given via a free lumen. However, if required IV compatibility includes: calcium gluconate, fentanyl, furosemide (in sodium chloride only), magnesium sulfate, , morphine, potassium chloride (in sodium chloride only),
IV Line incompatibility	Vancomycin

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS See Summary of Product Characteristics and BNF for Children

FURTHER INFORMATION

- Nebulised salbutamol 2.5-5 mg may be used to treat hyperkalaemia, although IV injection is preferred.
- May cause tachycardia, tremor and lactic acidosis. This may worsen work of breathing.
- Give centrally where possible, otherwise via a large peripheral vein.

LICENSED STATUS Indication is unlicensed and IV route is not licensed for children under 12 years

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Neonatal

Document Number:	004	Supersedes:	003
Prepared By/Checked By	Andrew Stockton/Lauren Williams/Olivia Scales	Final Approval By	WoS Neonatal Pharmacist Group
Date Prepared	September 2026	Review Date	September 2029

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Network. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department