Oxybutynin

****This monograph is under review****

FORM Liquid 2.5mg/5ml

INDICATION Neurogenic bladder instability

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Neonate to six months	100 – 200 micrograms/kg	2 -3 times daily	Oral

Maximum dose – 1.25mg THREE times daily

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

STORAGE

LICENSED STATUS Unlicensed dosage

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared by:	Peter Mulholland	Checked by	WoS Neonatal Pharmacists
Date prepared	January 2017	Date updated	October 2021
Updated by	Peter Mulholland	Review Date	October 2024
Extension Agreed By	WoS NPG	Review Date	April 2027

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist network. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.