Sodium Benzoate

This monograph is under review

Neonatal

FORM Ampoule containing 1g in 5ml (STOCKED IN RHC NICU)

INDICATION Treatment of hyperammonaemia

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
0 – 6 months	250mg/kg	Loading dose	IV
	Followed by 10mg/kg/hour Adjusted according to response	Continuous infusion.	IV

PRESCRIPTION OF CONTINUOUS INFUSION

1g in 20ml infusion fluid
This gives;

- 10mg/kg/hr = 0.2ml/kg/hr

RECONSTITUTION

Already in solution

DILUTION

Sodium Benzoate 200mg/ml	5ml
Glucose 10%	Up to 20ml total

Gives a 50mg in 1ml solution. Use the required volume.

METHOD OF ADMINISTRATION

Loading Dose
5 x weight (kg) = number of mL of diluted solution (50mg/ml) to be infused over 90 minutes

Continuous Infusion
By continuous intravenous infusion, flow rate adjusted according to response.

COMPATIBILITY

Solution compatibility	Sodium chloride 0.9% and 0.45%, glucose 5% and 10%.
Solution incompatibility	No information
IV Line compatibility	Can be infused at Y-site with arginine, carnitine and sodium phenylbuytrate (no data to support this but has been done in practice)
IV Line incompatibility	No information

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION Undiluted injection contains 1.4mmol Sodium per ml

NEONATAL West of Scotland Drug Monographs
Parenteral Route

LICENSED STATUS Unlicensed.

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared by:	WOS Neo Pharm Group	Checked by	Anisa Patel
Date prepared	May 2016	Date updated	June 2018
Updated by	Hazel Fisher	Review Date	June 2021
Updated by	Anisa Patel	Date updated	November 2020
Review date	November 2023		
Extension Agreed By	WoS NPG	Review Date	August 2027

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Network. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department