Folic Acid

****This monograph is under review****

FORM Oral Solution 2.5mg/5ml

INDICATION Prevention of late anaemia in haemolytic disease of the newborn

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Preterm	500microgram	ONCE daily	Oral
Term	1mg	ONCE daily	Oral

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children (links below)

LICENSED STATUS Licensed for use in all ages

LINKS BNF for Children: / Electronic Medicines Compendium

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared by:	WoS Neo Pharm Group	Checked by	Peter Mulholland
Date prepared	August 2016	Date updated	October 2021
Updated by	Peter Mulholland	Review Date	October 2024
Extension Agreed By	WoS NPG	Review Date	January 2027

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the package insert, BNF for Children, and Summary of Product Characteristics. For further advice contact your clinical pharmacist or pharmacy department.