Teicoplanin

****This monograph is under review****

FORM Vial containing 200mg powder for reconstitution

INDICATION Glycopeptide antibiotic for the management of serious gram-positive infections

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Neonate	16mg/kg loading dose	ONE DOSE ONLY	IV infusion
	Then 8mg/kg	ONCE daily Start 24 hours after loading dose	IV infusion
1 – 6 months	Initially 10 mg/kg	Every 12 hours for 3 doses	IV infusion
	Then 10mg/kg	ONCE daily	IV infusion

RECONSTITUTION Reconstitute each vial with the accompanying ampoule of water for injection (3.2ml) and roll vial **gently** until completely dissolved. If foaming occurs, leave standing for about 15 minutes to allow foam to subside. This will provide a solution of 67mg/ml (200mg vial), as excess is included in the vial.

DILUTION

Teicoplanin 200mg/3ml	3ml
Sodium Chloride 0.9% inj	Up to 20ml total

Gives a 10mg in 1ml solution. Use the required volume.

METHOD OF ADMINISTRATION IV Infusion (preferred route in neonates):
Administer over 30 minutes

If clinically necessary can be given as a slow IV bolus injection over 3-5 minutes.

COMPATIBILITY

Solution compatibility	Sodium chloride 0.9%, Glucose 5%; Sodium Chloride 0.45% and Glucose 5%
Solution incompatibility	No information
IV Line compatibility	No information
IV Line incompatibility	No information

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

NEONATAL West of Scotland Drug Monographs
Parenteral Route

- See Summary of Product Characteristics and most recent edition of BNF for Children

PH 7.5

LICENSED STATUS **Licensed in all ages**

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	001	Supersedes:	Nil
Prepared by:	WOS Neo Pharm Group	Checked by	Anisa Patel
Date prepared	May 2016	Date updated	September 2019
Updated by	A Patel	Review Date	September 2022
Extension Agreed By	WoS NPG	Review Date	July 2027

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Network. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department