Atropine

****This monograph is under review****

Neonatal

FORM

Ampoule containing 600 micrograms in 1 ml

NB. Please note higher strength of this product. Further dilution required

INDICATION As premedication, use if required.

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth to 6 months	15 microgram/kg	Single dose	I.V.bolus over a few seconds, followed by a 0.5-1ml flush over a few seconds

RECONSTITUTION Already in solution

DILUTION

Atropine Injection 600micrograms/ml	0.5ml
Sodium Chloride 0.9% injection	up to 10ml total

Gives a 30microgram in 1ml solution. Use the required volume.

Dosage guide:

- 0.5ml per kg = 15microgram/kg

METHOD OF ADMINISTRATION I.V. injection over a few seconds followed by a 0.5-1ml flush over a few seconds

COMPATIBILITY

Solution compatibility	Sodium chloride 0.9%.
Solution incompatibility	No information.
IV Line compatibility	
IV Line incompatibility	

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION Atropine given intravenously should produce an effect within 30 seconds and may last approximately 2 to 6 hours.

PH 2.8 – 4.5

LICENSED STATUS Not licensed for this indication in children under 12 years

APPLICABLE POLICIES West of Scotland Neonatal Guidelines
Consult local policy if applicable

Document Number:	004	Supersedes:	003
Prepared by:	Maria Tracy	Checked by	WoS Neonatal Pharmacists
Date prepared	October 2023	Review Date	October 2026
Extension Agreed By:	WoS NPG	Review Date	April 2027

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Network. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department