Domperidone

****This monograph is under review****

FORM Oral Suspension 1mg/ml

INDICATION Management of Gastro-oesophageal reflux disease (GORD)

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth to 6 months	250micrograms/kg*	EIGHT hourly	Oral

*If symptoms are uncontrolled on above dose it can be increased to a maximum of 400microgram/kg three times daily. This is a consultant decision and regular cardiac monitoring is recommended.

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

- See Summary of Product Characteristics and most recent edition of BNF for Children
  - Domperidone is associated with a small increased risk of cardiac side effects including QT interval prolongation and ventricular arrhythmias
  - An ECG should be performed before commencing treatment and before any dose change

FURTHER INFORMATION

- The effect of domperidone on gastro-intestinal activity is antagonised by morphine and other opioid analgesics

LICENSED STATUS Unlicensed Use

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Neonatal

Document Number:	002	Supersedes:	001
Prepared by:	Wos Neo Pharm Group	Checked by	Marie Stewart
Date prepared	Aug 2016	Date updated	October 2023
Updated by	P. Mulholland/ L. Williams	Review Date	October 2026
Extension Agreed By	WoS NPG	Review Date	October 2027

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist network.
Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.