Chlorothiazide

****This monograph is under review****

FORM 250mg/5ml oral suspension

INDICATION
1. Diuresis in the management of Bronchopulmonary Dysplasia (BPD)
2. Hyperinsulinism (in combination with Diazoxide)

DOSE RANGE

1. Bronchopulmonary Dysplasia

AGE	DOSE	FREQUENCY	ROUTE
Birth – 6 months	10-20mg/kg	TWICE Daily	Oral

2. Hyperinsulinism

AGE	DOSE	FREQUENCY	ROUTE
Birth – 6 months	3-5mg/kg	TWICE Daily	Oral

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

- See Summary of Product Characteristics and most recent edition of BNF for Children
  - Can cause electrolyte disturbance. Monitor sodium and potassium level.

LICENSED STATUS Unlicensed medicine

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Neonatal

Document Number:	002	Supersedes:	001
Prepared by:	WoS Neo pharm group	Checked by	Maria Tracey
Date prepared	Aug 2016	Date updated	August 2021
Updated by	M. Tracey	Review Date	August 2024
Extension Agreed By	WoS NPG	Review Date	May 2027

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist network. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.