Diazoxide

****This monograph is under review****

FORM Oral suspension 50mg/ml

INDICATION Hyperinsulinism

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
0-6 months	Initially 1.7mg/kg THREE times daily to establish a response Adjust dose according to response. Usual maintenance dose 1.5mg-3mg/kg*	2 -3 times daily	Oral

* Up to 7mg/kg 3 times daily may be required in some cases; higher doses unlikely to be beneficial.

- **Total fluid volume should be restricted to 130ml/kg/day for at least 24 hours prior to starting diazoxide treatment, whilst maintaining euglycaemia.**
- **Commence treatment with chlorothiazide in line with the neonatal monograph.**

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

LICENSED STATUS Unlicensed medicine

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared by:	WoS Neo Pharm Group	Checked by	Peter Mulholland
Date prepared	August 2016	Date updated	May 2020
Updated by	Anisa Patel	Review Date	May 2023
Extension Agreed By	WoS NPG	Review Date	January 2027

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist network. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.