SODIUM FEREDETATE (IRON)

FORM Oral Solution
Contains sodium feredetate 190mg (equivalent to 27.5mg of iron) in 5mls

INDICATION Prevention or treatment of iron deficient anaemia

DOSE RANGE

1. PROPHYLAXIS - premature infants less than 34 weeks gestation from 28 days of age or on discharge from unit if sooner, until ONE year of age.

AGE	DOSE	FREQUENCY	ROUTE
Premature Infants (see local policy)	0.5 - 1ml	Once daily	Oral

Dose may be split (0.5ml twice daily) to improve tolerability.
Consider dose reduction if patient is receiving SMA fortifier as this contains iron.

2 TREATMENT

AGE	DOSE	FREQUENCY	ROUTE
Neonate	1.5ml (reduced if not tolerated to 1.5ml once daily on alternate days)	Once daily	Oral
1 – 6 months	2ml (reduced if not tolerated to 2ml once daily on alternate days)	Once daily	Oral

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

- Magnesium, calcium and zinc salts may reduce the absorption of iron preparations therefore, where possible, these medicines should be administered at different times.

LICENSED STATUS Not licensed for prophylaxis in preterm infants.

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Neonatal

NEONATAL West of Scotland Drug Monographs
ORAL/OTHER ROUTES

Document Number:	003	Supersedes:	002
Prepared By/Checked By	Jacqueline Rooney / Frances Lowrie/Marie Stewart	Final Approval By	WoS Neonatal Pharmacist Group
Date Prepared	August 2026	Review Date	August 2029

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.