Protamine Sulfate

****This monograph is under review****

Neonatal

FORM Ampoule containing 50mg in 5ml

INDICATION Heparin Antidote

DOSE RANGE

AGE	TIME SINCE HEPARIN GIVEN (minutes)	PROTAMINE DOSE PER 100units OF HEPARIN	ROUTE
Neonate – 6 months	<30	1mg	IV
	30-60	0.5 -0.75mg	IV
	60-120	0.375 – 0.5mg	IV
	>120	0.25 – 0.375mg	IV

Maximum 50mg (5ml) per dose

RECONSTITUTION Already in solution

DILUTION Not required

METHOD OF ADMINISTRATION IV injection over 10minutes undiluted, Dilute if required with sodium chloride 0.9%

COMPATIBILITY

Solution compatibility	Sodium chloride 0.9%
Solution incompatibility	
IV Line compatibility	None
IV Line incompatibility	All other medicines

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

SPECIAL MONITORING REQUIREMENTS Monitor APTT and INR

FURTHER INFORMATION In excess protamine has an anticoagulant effect

PH 2.5 – 3.5

LICENSED STATUS Unlicensed in neonates

APPLICABLE POLICIES West of Scotland Neonatal Guidelines
Consult local policy if applicable

Document Number:	001	Supersedes:	Nil
Prepared by:	Peter Mulholland	Checked by	Lauren Williams
Date prepared	June 2018	Date updated	
Updated by		Review Date	June 2021
Extension Agreed By	WoS NPG	Review Date	May 2027

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Network. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department