Levetiracetam

FORM Vial containing 500 mg in 5 ml

INDICATION Treatment of neonatal seizures

Neonatal

LOADING DOSE

AGE	DOSE	FREQUENCY	ROUTE
All ages	**40 mg/kg** Followed by additional STAT doses of 10mg/kg if required. Max total loading dose 60mg/kg	STAT loading dose	IV

STARTING MAINTENANCE DOSE

GESTATIONAL AGE	AGE	DOSE	FREQUENCY	ROUTE
<36 weeks	< 7 days	5 – 10 mg/kg*	Every 8 hours	IV
	≥ 7 days	10 – 20 mg/kg*	Every 12 hours	IV

GESTATIONAL AGE	AGE	DOSE	FREQUENCY	ROUTE
≥ 36 weeks	< 7 days	7.5 – 10 mg/kg*	Every 8 hours	IV
	≥ 7 days	10 – 30 mg/kg*	Every 12 hours	IV

*Starting doses/standard dosing ranges only. Doses up to 60mg/kg/day (pre-term infants) and 100mg/kg (term infants) have been given in some studies (on specialist Neurology advice only).

CAUTION in renal impairment – maintenance dose adjustments required. Discuss with Pharmacy.

RECONSTITUTION Already in solution

DILUTION

Levetiracetam injection 500 mg/5 ml	2 ml
Sodium chloride 0.9%	Up to 20 ml total

Gives a 10mg/ml solution. Use the required volume.

METHOD OF ADMINISTRATION Slow IV infusion over 15 minutes

COMPATIBILITY

Solution compatibility	Sodium chloride 0.9%, Glucose 5%
Solution incompatibility	No information
IV Line compatibility	Diazepam, lorazepam
IV Line incompatibility	No information

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

LICENSED STATUS Not licensed for monotherapy in neonates

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared by/Checked by:	Susan Kafka/Lauren Williams/Andrew Stockton	Approved by:	WoS NPG
Date prepared	October 2017	Review Date	August 2029

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department