# Acetylcysteine

****This monograph is under review****

**FORM** Ampoule containing 2g in 10ml (20%)

**INDICATION** Mucolytic adjuvant in the therapy of respiratory disorders associated with thick, viscous, mucus hypersecretion.
Fistula / Rectal washout

**DOSE RANGE**

| AGE | DOSE | FREQUENCY | ROUTE |
|---|---|---|---|
| **MUCOLYTIC** | | | |
| Birth to 6 months | 40 - 60mg/hour (1 - 1.5ml/hour) | Continuous | NasoGastric |
| | | | |
| **FISTULA WASHOUT** | | | |
| Birth to 6 months | 5ml of 4% solution | Every TWELVE hours | Via fistula |
| | | | |
| **RECTAL WASHOUT** | | | |
| Birth to 6 months | 20ml of 4% solution | ONCE daily | Rectal |

Neonatal

**CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS**
- See Summary of Product Characteristics and most recent edition of BNF for Children

**FURTHER INFORMATION** Dilute before use to give a 4% solution

| | |
|---|---|
| Acetylcysteine injection 2g/10ml | 4ml |
| Sodium chloride 0.9% | to 20ml |

**STORAGE** Use immediately

**LICENSED STATUS** Unlicensed

**APPLICABLE POLICIES** West of Scotland Neonatal Guidelines:

Consult local policy if applicable

| | | | |
|---|---|---|---|
| Document Number: | 001 | Supersedes: | Nil |
| Prepared by: | Peter Mulholland | Checked by | WoS Neonatal Pharmacists |
| Date prepared | February 2020 | Date updated | |
| Updated by | | Review Date | February 2023 |
| Extension Agreed By | WoS NPG | Review date | March 2027 |

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist network. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.