Trimethoprim

FORM Oral suspension 50mg/5ml

INDICATION

1. Prophylaxis in renal problems
2. Cover for micturating cystourethrogram (MCUG)
3. Treatment of susceptible infections

DOSE RANGE

1. Prophylaxis

AGE	DOSE	FREQUENCY	ROUTE
Birth to 6 months	2mg/kg	ONCE daily at night	Oral

2. MCUG Cover

AGE	DOSE	FREQUENCY	ROUTE
Birth to 6 months	2mg/kg	Every 12 hours for 72 hours *To be started on day before planned MCUG procedure*	Oral

3. Treatment of Susceptible Infections

AGE	DOSE	FREQUENCY	ROUTE
0 – 28 days	3mg/kg as a STAT dose Then 1-2mg/kg/dose	 Every 12 hours	Oral
1-6 months	4mg/kg/dose	Every 12 hours	Oral

FURTHER INFORMATION There is a theoretical risk of folate deficiency with long term use in neonates. Calcium folinate supplements may be needed.

Doses should be reduced in severe renal impairment – contact pharmacy for dosing advice

STORAGE Store in original packaging below 25^{O}C

LICENSED STATUS Not licensed in children under 6 weeks

Not licensed for use in children under 6 months for the prophylaxis of recurrent urinary tract infection

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Neonatal

NEONATAL West of Scotland Drug Monographs
ORAL/OTHER ROUTES

Document Number:	002	Supersedes:	001
Prepared By/Checked By	Lauren Williams/Julie Lowther/Frances Lowrie/Susan Kafka	Final Approval By	WoS Neonatal Pharmacist Group
Date Prepared	September 2026	Review Date	September 2027

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.