PARACETAMOL - Analgesia / Pyrexia

Neonatal

FORM Ampoule containing 100mg in 10ml

INDICATION Analgesic for short-term treatment of moderate pain or fever

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
<32 weeks CGA	7.5mg/kg/dose	12hourly	IV
32 weeks – 1month CGA	7.5mg/kg/dose	8 hourly	IV
1month CGA up to 10kg	7.5mg – 10mg/kg/dose (max 30mg/kg/day)	6 - 8hourly	IV

RECONSTITUTION Already in solution

DILUTION Can be administered undiluted

METHOD OF ADMINISTRATION Administer as an intravenous infusion over 15 minutes

COMPATIBILITY

Solution compatibility	Sodium chloride 0.9%, Glucose 5%
Solution incompatibility	All other IV fluids
IV Line compatibility	Caffeine citrate, Cefotaxime, Dexamethasone, Esmolol, Gentamicin, Hydrocortisone succinate, Meropenem, Midazolam, Morphine, Piperacillin with Tazobactam, Vancomycin, Parenteral Nutrition including lipid
IV Line incompatibility	All other drugs

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
See Summary of Product Characteristics and most recent edition of BNF for Children.
Caution in hepatic impairment or neonates with unconjugated hyperbilirubinaemia – monitor hepatic function

FURTHER INFORMATION

- Do not give with any other preparation containing paracetamol
- Doses higher than the recommended entail the risk of serious liver damage. Clinical signs and symptoms of liver damage are not usually seen until 2-6 days after administration

PH 5.5

STORAGE Room temperature

NEONATAL West of Scotland Drug Monographs
Parenteral Routes

LICENSED STATUS Not licensed in neonates or children under 10kg

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	004	Supersedes:	003
Prepared By/Checked By	Julie Lowther/Olivia Scales/Natalie Donald	Final Approval By	WoS Neonatal Pharmacist Group
Date Prepared	August 2026	Review Date	August 2029

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department