Carnitine (Levocarnitine)

****This monograph is under review****

Neonatal

FORM Ampoule containing 1g in 5ml

INDICATION Treatment of primary and secondary carnitine deficiency
N.B. Consultant Paediatrician must authorise use.

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth to 6 months	*initially 25mg/kg/dose	every 6 hours	IV
	may increase to 175mg/kg/day **or** 3g/day		

* Specialist consultants may recommend different doses on an individual baby basis.

RECONSTITUTION Already in solution

DILUTION (if required)

L-Carnitine 200mg/ml inj (Carnitor®)	5ml
Glucose 5%	to total 10ml

This gives 100mg/ml Carnitine solution. Use the required volume.

METHOD OF ADMINISTRATION Slow IV injection over 2 – 3 minutes.

COMPATIBILITY

Solution compatibility	Glucose 5%, Glucose 10%, sodium chloride 0.9%, Ringers solution
Solution incompatibility	No information
IV Line compatibility	Infuse at y-site with arginine, sodium benzoate and sodium phenylbutyrate
IV Line incompatibility	No information

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION Extravasation risk.

PH 6.0 to 6.5

LICENSED STATUS Injection licensed in neonates and children. Single dose vial only licensed for patients > 12 years

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	002	Supersedes:	Nil
Prepared by:	WoS Neo Pharm Group	Checked by	Marie L Currie
Date prepared	May 2016	Date updated	March 2018
Updated by	Hazel Fisher	Review Date	March 2021
Extension Agreed by:	WoS NPG	Review Date	August 2027

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Network. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department