Nystatin

FORM Oral Suspension 100 000unit/ml

INDICATION Treatment of Oral and GI tract candida infections

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth to 6 months	1ml (100 000unit)	4 times daily after feeds usually for 7 days, continue for 48 hours after lesions heal	Oral

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

- Oral medicine to be dropped into the mouth after feeds. The longer the suspension is kept in contact with the affected area in the mouth, before swallowing, the greater will be its effect.
- Treatment of perineal thrush usually requires combined topical and oral therapy to eliminate candida from GI tract.

LICENSED STATUS Not licensed in neonates

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared by/ Checked by:	Jacqueline Rooney/Frances Lowrie/Julie Lowther	Final approval by:	WoS Neonatal Pharmacist Group
Date prepared	September 2026	Review date:	September 2029

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.