Omeprazole

FORM

Omeprazole 20mg/5ml oral solution (ACLOMEP)
Omeprazole 10mg dispersible tablets (Losec MUPS)

Caution: there are other available formulations that are not recommended for use

INDICATION

Gastro-oesophageal reflux disease

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth – 6 months	700 micrograms/kg-3mg/kg Maximum dose 20mg	ONCE daily	Oral

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

- See Summary of Product Characteristics and most recent edition of BNF for Children (links below)

FURTHER INFORMATION

- **Consult local guidance for appropriate choice of preparation**
- Losec MUPS tablets can be halved and doses rounded to the nearest 5mg, where appropriate. Each tablet (or half tablet) should be dispersed in 5-10ml of sterile water. Do not crush the tablet or manipulate further than half.
- Omeprazole should, where possible, be given approximately one hour before feeds on an empty stomach. In practice it has been accepted that mixing with a small volume of milk for administration improves tolerability.

STORAGE

- Once reconstituted, ACLOMEP liquid should be stored in the fridge and used within after 28 days.
- Losec MUPS tablets may take 5-10 minutes to fully disperse.

LICENSED STATUS

- ACLOMEP liquid – unlicensed medication
- Losec MUPS-off label use of licensed product

APPLICABLE POLICIES

West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	004	Supersedes:	003
Prepared by:	Julie Lowther/Jacqueline Rooney/Frances Lowrie	Final Approval By	WoS Neonatal Pharmacist Group
Date Prepared	September 2026	Date updated	September 2029

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.

Neonatal