Isoprenaline

****This monograph is under review****

FORM Ampoule containing 1 mg in 5ml

INDICATION Complete heart block

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
0 - 6 months	0.1 – 1 microgram/kg/minute	Continuous Infusion	IV

PRESCRIPTION OF CONTINUOUS INFUSION

600 micrograms/kg of isoprenaline in 50ml of infusion fluid

This gives:

- 0.1 micrograms/kg/min at 0.5ml/hr

RECONSTITUTION Already in solution

DILUTION

For continuous Infusion

Using 1 mg/5ml injection

3 x wt (kg) is the number of ml of 1 mg/5ml to be diluted up to 50ml total with infusion fluid (equivalent **600 micrograms/kg in 50ml**)

METHOD OF ADMINISTRATION By continuous intravenous infusion, flow rate adjusted according to the baby's response (see prescription section for details).

COMPATIBILITY

Solution compatibility	Glucose 5%, Sodium Chloride 0.9%
Solution incompatibility	No information
IV Line compatibility	dobutamine, dopamine, heparin, milrinone (glucose only), noradrenaline, potassium chloride, sodium nitoprusside (glucose only), TPN, vecuronium (glucose only)
IV Line incompatibility	All other drugs

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

See Summary of Product Characteristics and most recent edition of BNF for Children

.

SPECIAL MONITORING REQUIREMENTS ECG, arterial blood pressure, heart rate, urine flow, central venous pressure, blood pH, blood pCO_2 or bicarbonate, and cardiac output.

FURTHER INFORMATION Discard the injection if it is pinkish or darker than slightly yellow or contains a precipitate.

SODIUM CONTENT

PH

LICENSED STATUS Licensed product

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared by:	WoS Neo Pharm group	Checked by	Peter Mulholland
Date prepared	May 2016	Date updated	September 2022
Updated by	Anisa Patel	Review Date	September 2024
Extension Agreed By	WoS NPG	Review Date	March 2027

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Network. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department