Calcium Resonium® - Rectal

CAUTION: this monograph is for Calcium Resonium (Sanofi). There are multiple brands of calcium polystyrene sulfonate, follow the manufacturer's directions of available brand.

BRAND/ MANUFACTURER Calcium Resonium® (Sanofi)

FORM **Calcium Resonium®** powder for oral/rectal use (contains calcium polystyrene sulfonate 99.9%w/w)

INDICATION Management of Hyperkalaemia

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth – 6 months	500mg – 1 gram/kg/dose	Once daily	**Rectal**

METHOD OF ADMINISTRATION [1,2,3]

Calcium Resonium® Mix ONE scoop (15gram) of resin with 75ml of sterile water or glucose 10% and administer immediately.

Resulting Solution
500mg/kg = 2.5ml/kg
1g/kg = 5ml/kg

Administer required dose immediately
Irrigate colon to remove resin after 8-12 hours.

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

- Contraindicated in neonates with reduced gut motility and bowel obstruction.
- Due to the risk of digestive haemorrhage or colonic necrosis, particular care should be observed in premature infants or low birth weight infants

FURTHER INFORMATION

- Calcium is released in the gut; babies should be monitored for evidence of hypercalcaemia.
- Exchanges 1-2mmol of potassium per gram of resin.
- Calcium Resonium® should be stopped when serum potassium is below 5mmol/L

LICENSED STATUS Licensed Use

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Neonatal

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.