ABIDEC®

FORM Oral drops

INDICATION Vitamin supplementation of premature neonates <34 weeks

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Premature infants <34 weeks	0.6ml	ONCE daily*	Oral

* If an infant persistently vomits following 0.6ml Abidec® once daily this can be given as 0.3ml twice daily to improve tolerability.

*Consider a dose reduction if patient is also receiving fortifier as per local protocols

Commence as soon as enteral feeding has been established (at least 100ml/kg/day tolerated).

Abidec® should be started for babies weaning from TPN once enteral feeding (at least 100ml/kg/day) is tolerated, regardless of ongoing provision of SMOFvits®

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION 0.6ml contains Vitamin A 1333units
Thiamine 0.4mg
Riboflavin 0.8mg
Pyridoxine 0.8mg
Nicotinamide 8mg
Vitamin C 40mg
Vitamin D 400units

Abidec® contains refined arachis (peanut) oil

STORAGE Store at room temperature. Discard 4 weeks after opening

LICENSED STATUS Licensed use. Dose for premature babies whilst in hospital is off label but endorsed in BNFC.

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared By/Checked By	Jacqueline Rooney / Frances Lowrie/Andrew Stockton	Final Approval By	WoS Neonatal Pharmacist Group
Date Prepared	August 2026	Review Date	August 2029

NEONATAL West of Scotland Drug Monographs
ORAL/OTHER ROUTES

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.