Furosemide

FORM 20mg/5ml oral solution
Special care required – other strengths available

INDICATION Diuresis in cardiac or renal failure, pulmonary oedema or fluid overload.

Neonatal

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Neonate up to 31 weeks CGA	500microgram to 2mg/kg	Once daily	Oral
Neonate >31 weeks CGA	500microgram to 2mg/kg	1-2 times daily	Oral
Infant 1– 6 months	500microgram to 2mg/kg Higher doses may be required in resistant oedema. Maximum 12mg/kg per day Maximum daily dose 80mg per day	2-3 times daily	Oral

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

- Furosemide oral solution contains ethanol as an excipient with the quantity varying between brands. Caution may be required in larger doses. Further information can be provided by pharmacy
- A marked fall in blood pressure may occur with concomitant treatment with an ACE inhibitor
- Nephrocalcinosis has been reported in premature neonates
- Monitor electrolytes and fluid balance during treatment

LICENSED STATUS Licensed product

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared by:	Olivia Scales/F Lowrie/ J Lowther	Final Approval BY	WoS Neonatal Pharmacist Group
Date Prepared	September 2026	Review Date	September 2029

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.