Midazolam

This monograph is under review

DISCUSS WITH CONSULTANT NEONATOLOGIST BEFORE USE

FORM Oromucosal solution midazolam 10mg/ml

INDICATION Control of Seizures

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Neonate – 3 months	300micrograms/kg (max 2.5mg)	Single dose. Repeated ONCE after 10 minutes if necessary	Buccal
3-6 months	2.5mg		

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

- Administer half of the required dose quickly to each side of the buccal cavity (between lower cheeks and gums)
- Initial effect should be seen after 5 minutes
- Reduce dose in liver failure
- Flumazenil is the antidote for midazolam induced respiratory depression
- Erythromycin and possibly fluconazole may inhibit the metabolism of midazolam, producing profound sedation.

STORAGE Store at room temperature. Cap must be replaced immediately otherwise crystals of sweetener may form. Discard if solution not clear.

LICENSED STATUS Unlicensed product

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared by:	WoS Neo pharm group	Checked by	June Grant
Date prepared	Aug 2016	Date updated	August 2021
Updated by	Roisin McCabe	Review Date	August 2024
Extension Agreed By	WoS NPG	Review Date	February 2027

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist network. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.