NEONATAL RESUSCITATION DRUGS				
DRUG	**DOSE**	**DILUTION INSTRUCTIONS**	**ADMINISTRATION**	**COMMENTS**
ADRENALINE (EPINEPHRINE) **1:10,000** **100microgram/ml**	**20microgram/kg**	No need to dilute Draw up required volume into a 1ml syringe	**0.2ml x weight (kg) of 1:10,000 strength** Rapid IV bolus / UVC / intraosseous	Repeat dose every 3-5 minutes if required and consider use of sodium bicarbonate See Adrenaline(for resuscitation) monograph for endotracheal (ET) dose
SODIUM BICARBONATE 4.2%	**2ml/kg**	**No need to dilute**	**Give 2ml X weight (kg)** Slow IV, or via UVC over 3-5 min.	Single dose May be repeated **Incompatible with all other drugs and fluids**
GLUCOSE 10%	**2.5ml/kg**	**No need to dilute**	**IV** Over 10-20 seconds	
Sodium Chloride 0.9%	**10ml/kg**	**No need to dilute**	**IV** over 10-20 seconds	
Naloxone				
NB Naloxone is NOT a resuscitation drug. It may be required, to reverse opiate induced sedation, following resuscitation				
NALOXONE (NARCAN) 400microgram/ml	**200micrograms**	**No need to dilute**	**Give 0.5ml IM into the anterior thigh once only**	Reversal of narcotic-induced respiratory depression following maternal use in labour. Do not give to infant whose mother is an opiate user
Vitamin K at birth doses:				
VITAMIN K Phytomenadione (Konakion) **2mg in 0.2ml**	**Gestation < 36 $^{+0}$ weeks Dose= 0.5mg IM**		**Gestation ≥ 36 $^{+0}$ weeks Dose= 1mg IM**	

NEONATAL INTUBATION DRUGS

N.B. For Hypoplastic Left Heart / complex cardiac patients – See separate local guidance

DRUG	DOSE	DILUTION INSTRUCTIONS	ADMINISTRATION	COMMENTS
ATROPINE 600microgram/ml	**15microgram/kg**	**Dilute 300micrograms (0.5ml) to 10ml with 0.9% saline = 30microgram in 1ml**	**Give 0.5ml X weight (kg) of 30microgram/ml solution**	IV injection over a few seconds., followed by a 05-1ml flush over a few seconds
FENTANYL Intubation kit 10microgram/ml	**5microgram/kg**	**No need to dilute**	**Give 0.5ml X weight (kg)** of 10microgram/ml solution	Single dose, slow IV injection , over 60 seconds followed by a 0.5-1ml flush over 60 seconds
FENTANYL 100microgram/2ml If no kits available	**5microgram/kg**	**Dilute 25 micrograms (0.5ml) to 2.5ml total with 0.9% saline = 10microgram in 1ml**	**Give 0.5ml X weight (kg) of 10microgram/ml solution**	Single dose, slow IV injection , over 60 seconds followed by a 0.5-1ml flush over 60 seconds
SUXAMETHONIUM Intubation kit 10mg in 2.5ml	**2mg/kg**	**No need to dilute**	**Give 0.5ml X weight (kg)** of 10mg in 2.5ml solution	Single dose, IV injection over a few seconds, followed by a 0.5-1ml flush over a few seconds Repeated once if necessary after 2 – 6 minutes
SUXAMETHONIUM 100mg / 2ml If no kits available	**2mg/kg**	**Dilute 10mg (0.2ml) to 2.5ml with 0.9% saline = 4mg in 1ml (prepare 2 doses)**	**Give 0.5ml X weight of 4mg/ml solution**	Single dose, IV injection over a few seconds, followed by a 0.5-1ml flush over a few seconds Repeated once if necessary after 2 – 6 minutes
MORPHINE 10mg / ml	**100 microgram/kg**	**Dilute 4mg (0.4ml) to 20ml with 0.9% saline = 200microgram in 1ml**	**Give 0.5ml X weight (kg)** of **200microgram in 1ml solution**	Single dose IV injection over 5min Repeated once if necessary.

Document Number: 004 Supersedes: 003 Date prepared: April 2018 Review date: Oct 2026
Prepared by: GG&C Neonatal Pharmacists Checked by: Maria Tracey Updated by: M Tracey Oct 2023
Extension Agreed by: WoS NPG Review Date: April 2027