Amlodipine

FORM Oral liquid 1mg/1ml

INDICATION Hypertension

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth – 6 months	100 – 200 micrograms/kg	ONCE daily	Oral

*Dose increased if necessary up to 400 micrograms/kg once daily, according to response (usually at 1-2 week intervals).
** Total daily dose may be split twice daily if required

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION
Dose may need to be reduced in hepatic impairment
Product should **NOT** be shaken prior to administration
Use within 30 days of opening

STORAGE
Bottle should be stored in outer container to protect from light
Some products may require fridge storage – please check packaging

LICENSED STATUS Unlicensed in children under 6 years old

APPLICABLE POLICIES
West of Scotland Neonatal Guidelines
Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared b/Checked By	Lauren Williams/Frances Lowrie/Olivia Scales	Final Approval By	WoS Neonatal Pharmacist Group
Date Prepared	September 2026	Review Date	September 2029

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland pharmacist group. Information is correct at the time of publication and as per local practice agreement.
For further advice contact your clinical pharmacist or pharmacy department.