Amphotericin (Liposomal)

FORM Vial containing 50 mg powder for reconstitution

INDICATION Antifungal for the prophylaxis or treatment of severe systemic fungal infections

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Neonate	3 mg/kg (empiric) 5 mg/kg* (proven infection)	Once daily	IV infusion over 30 minutes
>1month	3 mg/kg (empiric) 5 mg/kg* (proven infection)	Once daily	IV infusion over 30 minutes

*Doses >5mg/kg may be recommended in some cases (e.g. proven mucormycoses infection) under the direction of the Infectious Diseases Team.

RECONSTITUTION Reconstitute each 50mg vial with 12ml of Water for Injection (to give final concentration of 4mg/ml). Shake vigorously for 30seconds.

DILUTION

Liposomal Amphotericin 4 mg/ml	5 ml (Draw up using the 5 micron filter provided)
Glucose 5%	up to 20 ml total

To give a 1 mg/ml solution then give the required volume.
(If a more concentrated solution is required, discuss with pharmacy)

METHOD OF ADMINISTRATION Infuse slowly over 30 minutes. Flush line before and after infusion with glucose 5% only.
(Doses >5 mg/kg should be infused over 2 hours)

COMPATIBILITY

Solution compatibility	Glucose 5%
Solution incompatibility	All other fluids
IV Line compatibility	Nil
IV Line incompatibility	All other drugs

THIS LIST IS NOT EXHAUSTIVE PLEASE CONTACT PHARMACY FOR FURTHER INFORMATION ON COMPATIBILITY WITH ANY MEDICINES NOT INCLUDED

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS
- See Summary of Product Characteristics and most recent edition of BNF for Children

FURTHER INFORMATION

PH 5.0-6.0

LICENSE STATUS Unlicensed in neonates.

NEONATAL West of Scotland Drug Monographs
Parenteral Route

APPLICABLE POLICIES West of Scotland Neonatal Guidelines

Consult local policy if applicable

Document Number:	004	Supersedes:	003
Prepared by/Checked by:	Susan Kafka/ J Rooney/A Stockton	Approved by	WoS Neonatal Pharmacist Group
Date prepared	June 2026	Review date	September 2029

Administer reconstituted solutions immediately.
All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications.
This monograph should be used in conjunction with the terms of reference document prepared by the West of Scotland Pharmacist Group. Information is correct at the time of publication and as per local practice agreement.
For further advice please contact your clinical pharmacist or pharmacy department